

of a ch

Right Eye 20/	Left Eye 20/	Pass	Fail

Right				Pass Fail
Left				Pass Fail

If your child does not attend pre-kindergarten or school away from the child care operation, one of the following must be presented when your child is admitted to the child care operation or within one week of admission. *(Select **only one** option.)*

- Health Care Professional's Statement: I have examined the above named child within the past year and find that he or she is abl

-

-

The following vaccines require multiple doses over time. Please provide the date your child received each dose.

Hepatitis B	Birth (first dose)	
	1–2 months (second dose)	
	6–18 months (third dose)	
Rotavirus	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
Diphtheria, Tetanus, Pertussis	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	15–18 months (fourth dose)	
	4–6 years (fifth dose)	
Haemophilus Influenza Type B	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	12–15 months (fourth dose)	
Pneumococcal	2 months (first dose)	
	4 months (second dose)	
	6 months (third dose)	
	12–15 months (fourth dose)	
Inactivated Poliovirus	2 months (first dose)	
	4 months (second dose)	
	6–18 months (third dose)	
	4–6 years (fourth dose)	
Influenza	Yearly, starting at 6 months. Two doses given at least four weeks apart are recommended for children who are getting the vaccine for the first time and for some other children in this age group.	
Measles, Mumps, Rubella	12–15 months (first dose)	
	4–6 years (second dose)	
Varicella	12–15 months (first dose)	
	4–6 years (second dose)	
Hepatitis A	12–23 months (first dose)	
	The second dose should be given 6 to 18 months after the first dose.	

Signature or stamp of physician or public health personnel verifying immunization information above:

Signature

Date Signed